

Effect of Some Plant Growth Regulators on Seed Firmness, Quantitative and Qualitative Characteristics of Pomegranate (*Punica granatum*) cv. Yousefkhani

S. Jafari, H. Sarikhani*, E. Ahmadi and S.M. Zahedi¹

The present study was carried out to investigate the effects of some plant growth regulators on quantitative and qualitative characteristics of pomegranate (*Punica granatum* cv. Yousefkhani) with emphasis on their effect on reducing seed firmness. In this study, 10 and 20 mg L⁻¹ kinetin, 10 and 40 mg L⁻¹ gibberellic acid, and paclobutrazol, as an anti-gibberellin, at four concentrations (10, 50, 100, and 200 mg L⁻¹) were sprayed four weeks after fruit set. Distilled water was applied as control. Results showed that foliar application of 10 and 20 mg L⁻¹ kinetin increased fruit length and fruit weight, respectively. In contrary, small fruits with lower edible portion were produced after foliar application of 200 mg L⁻¹. The highest percentage of edible portion was observed in fruits treated with 10 mg L⁻¹ kinetin plus 10 mg L⁻¹ gibberellic acid. Foliar application of plant growth regulators showed significant effects on seed firmness compared to control. The lowest seed firmness was observed using 100 mg L⁻¹ paclobutrazol. Foliar application with various plant growth regulators also affected the chemical properties of the fruits. While the other treatments reduced total anthocyanin levels, treatment by paclobutrazol at a concentration of 100 mg L⁻¹ preserved it.

Keywords: Gibberellic Acid, Kinetin, Paclobutrazol, Seed Firmness.

1. Former M.Sc. Student and Associate Professor, Department of Horticultural Science, Associate Professor, Department of Biosystem Engineering, Bu-Ali Sina University, Hamedan and Assistant Professor, Department of Horticultural Science, University of Maragheh, Maragheh, Iran, respectively.

* Corresponding author email: (sarikhani@basu.ac.ir).